

Full Length Research Paper

Biosorption of lead ions from aqueous solution by maize leaf

N. A. Adesola Babarinde^{1*}, J. Oyebamiji.Babalola² and R. Adebowale Sanni¹

¹Olabisi Onabanjo University, Ago-Iwoye, Nigeria.

²University of Ibadan, Ibadan, Nigeria.

Accepted 24 August, 2006

The removal of lead ions from dilute aqueous solution using maize (*Zea mays*) leaf as the adsorbent is reported in this paper. The effects of pH, initial metal ion concentration and contact time were studied at 27°C. The analysis of residual Pb(II) ions was determined using atomic absorption spectrophotometer. Maximum adsorption occurred at pH 3. The adsorption isotherms obtained at 27°C and optimum pH fitted well into both the Freundlich and Langmuir isotherms. The Freundlich and Langmuir equations are $\log \Gamma = 0.504 \log C_e + 0.6939$ and $1/\Gamma = 0.176/C_e - 0.03$, respectively. The correlation factors are 0.9959 and 0.9939. The result of the pH experiment shows that the initial pH would play a vital role in the removal of the lead ions from solution. The kinetic studies show that uptake of lead ions increases with time and that maximum adsorption was obtained within the first 30 min of the process. These results indicate that maize leaf has potential for removing lead ions from industrial wastewater.

Key words: Adsorption, Pb(II), Freundlich isotherm, Langmuir isotherm, maize leaf.

INTRODUCTION

The increase in industrial activity during recent years is greatly contributing to the increase of heavy metals in the environment, mainly in the aquatic systems (Marques et al., 2000). Water pollution due to heavy metals is an issue of great environmental concern (Vasudevan et al., 2003). Heavy metal ions such as cobalt, copper, nickel, chromium, lead and zinc are detected in the waste streams from mining operations, tanneries, electronics, electroplating, batteries and petrochemical industries as well as textile mill products. Major lead pollution is through automobiles and battery manufacturers.

Heavy metals have a harmful effect on human physiology and other biological systems when they exceed the tolerance levels (Koby et al., 2005). Exposure to lead can cause anaemia, diseases of the liver and kidneys, brain damage and ultimately death (Jain et al., 1989). Besides, chronic exposure to these contaminants present even at low concentrations in the environment can prove to be harmful to the human health. For these reasons, heavy metals must be removed

as much as possible from industrial effluents.

The conventional methods used to remove heavy metals include chemical precipitation, ion exchange, electrodialysis, membrane separations, reverse osmosis, and solvent extraction (Matlock et al., 2002; Feng et al., 2000; Mohammadi et al., 2004). The search for new, effective and economical technologies involving the removal of toxic metals from wastewaters has directed attention to biosorption based on metal binding capacities of various biological materials at little or no cost (Abia, et al., 2003; Ajmal et al., 2005; Aksu, 2002; Annadurai, et al., 2002; Babarinde and Oyedipe, 2001; Babarinde et al., 2002; Babarinde, 2002; Bansode, et al., 2003; Cimino, et al., 2000; Demirbas, 2003; Feng and Aldrich, 2004; Fourest and Roux, 1992). Of interest in this project work is the removal of lead ion from aqueous solution.

METHODS

Materials

The maize leaves (*Zea mays*) used as the adsorbent were obtained from a farm in Ijebu-Imusin, Ogun State, Nigeria. The dry leaves were rinsed with distilled water, sun-dried and cut into pieces. The salt used for the preparation of aqueous solutions was analytical grade.

*Corresponding authors E-mail: solababarinde@yahoo.com.
Tel: 2348037232934.

Biosorption experiments

Experiments were conducted at 27°C to study the effect of initial solution pH, ion concentration and contact time on the adsorption of lead ions. Each experiment was conducted in a thermostated water bath (Haake Wia model) and the residual metal ions analysed using Atomic Absorption Spectrophotometer (Buck Scientific model 210 VGP), at wavelengths of 283.3 nm for lead. The detection limit of 0.05 was used while air/acetylene mixture was used as the oxidant.

Effect of pH

The procedure used is similar to those earlier reported (Vasudevan et. al., 2003; Xu et. al., 2006). Experiments were conducted at 27°C to study the effect of initial solution pH on the adsorption of the metal ions by contacting 0.5 g of the maize leaf with 50 ml of 100 mg/l lead salt solution in a boiling tube. The pH of each of the metal solutions was adjusted to the desired value with 0.1 M sodium hydroxide and /or 0.1 M nitric acid (Lodeiro, et al., 2005).

The studies were conducted at pH values of 2, 3, 4, 5, 6 and 7. The boiling tubes containing the mixture were left in a water bath for 24 h. The sorbent was removed from the solution by filtration and the residual lead ion concentration in the solution was analysed by Atomic Absorption Spectrophotometer. All experiments were conducted in triplicates and the mean value was determined for each.

Effect of concentration

Batch adsorption study was carried out using a concentration range of 10–100 mg/l. 0.5 g of the maize leaf was weighed into each of the 30 boiling tubes employed and 50 ml of the metal ion solution was measured into each boiling tube containing the maize leaf. Three boiling tubes were used for a particular concentration. The adsorption mixture was then left in a water bath to maintain the temperature (27°C) for 24 h. The maize leaf was removed from the mixture by filtration and the concentration of the residual ions in the solution was determined using Atomic Absorption Spectrophotometer. The amount of metal ions adsorbed from solution was determined by difference. The mean value was also calculated.

The results obtained were analysed using both Freundlich (Freundlich, 1907) and Langmuir (Langmuir, 1918) isotherms. The Freundlich isotherm in linearised form is $\log \Gamma = 1/n \log C_e + \log K$, where K and n are Freundlich constants. The Langmuir isotherm in a linear form is $1/\Gamma = (1/b_m)(1/C_e) + 1/\Gamma_m$ where b_m is a coefficient related to the affinity between the sorbent and sorbate, and Γ_m is the maximum sorbate uptake under the given condition.

Effect of contact time

The adsorption of the metal ions by maize leaf was studied at various time intervals (0-120 min). A constant concentration of 100 mg/l was used. 0.5 g of maize leaf was weighed into thirty-six boiling tubes and 50 ml of each metal ion solution (100 mg/l) was added into each tube. The maize leaf in the thirty six tubes was then filtered at different time intervals of 5, 10, 15, 20, 25, 30, 45, 60, 75, 90, 105 and 120 min. The filtrates were then analyzed using Atomic Absorption Spectrophotometer. The amount of metal ions adsorbed was calculated for each sample. The mean of the three results for a particular time was then calculated and plotted against time.




Figure 1. Effect of pH on sorption behaviour of Pb(II) sorption by maize leaf.




Figure 2. The linearized Freundlich adsorption isotherm of Pb(II) using maize leaf.




Figure 3. The linearized Langmuir adsorption isotherm of Pb(II) using maize leaf.

Table 1. Freundlich and Langmuir isothermal adsorption parameters for the adsorption of Pb²⁺ ions at 27°C.

Freundlich parameters			Langmuir parameters		
1/n	K	R	1/b _m	1/Γ _m	R
0.504	4.356	0.9959	0.176	-0.03	0.9939




Figure 4. Time courses of Pb(II) ion removal by maize leaf.

RESULTS AND DISCUSSION

The results of the three different studies are shown in Figures 1 - 4. The result of the pH study shows that maximum sorption occurred at pH 3. The percentage Pb(II) adsorbed is slightly lower at higher pH. This suggests that optimum adsorption is obtained at pH 3. This implies that the initial pH would play a vital role in the removal of the Pb(II) ions from solution. The effect of metal ion concentration on adsorption capacity showed that maize leaf biomass adsorbed the lead ions from solution. The amount of Pb(II) ions adsorbed increased with increase in initial metal ion concentration. The data fitted well into both the Freundlich and Langmuir isotherms. The Freundlich and Langmuir parameters obtained (Table 1) compare well with those of other adsorbents that have been reported (Júnior, et al., 2003; Lodeiro, et al., 2005). The values of the parameters show that maize leaf is a good adsorbent for the removal of lead from wastewaters.

In conclusion, maize leaf, an agricultural waste, could be used as potential adsorbent for the removal of lead from aqueous solutions. The batch adsorption studies have shown that the adsorption of lead is pH dependent and the optimum pH for the removal is 3. The amount of lead ions adsorbed increased with increase in initial metal ion concentration. Maximum adsorption was obtained within the first 30 min.

ACKNOWLEDGEMENTS

The authors are grateful to J. O. A. Adegoke and T. Olayiwola of the Department of Chemical Sciences, Olabisi Onabanjo University, Ago-Iwoye for technical assistance.

REFERENCES

- Abia AA, Horsfall Jr. M, Didi O (2003). The use of chemically modified and unmodified cassava waste for the removal of Cd, Cu and Zn ions from aqueous solution. *Bioresour. Technol.* 90(3): 345–348.
- Ajmal M, Rao RAK, Anwar S, Ahmad J, Ahmad R (2003). Adsorption studies on rice husk: removal and recovery of Cd (II) from waste water. *Bioresour. Technol.* 86: 147–149.
- Aksu Z (2002). Determination of the equilibrium, kinetic and thermodynamic parameters of the batch biosorption of nickel (II) ion into *Chlorella*. *Process Biochem.* 38(1): 89–99.
- Annadurai G, Juang RS, Lee DJ (2002). Adsorption of heavy metals from water using banana and orange peels. *Water Sci-Technol.* 47(1): 185–190.
- Babarinde NAA, Oyedipe R A, (2001). Adsorption Characteristics of zinc (II) and cadmium (II) by Coconut husk and goat Hair. *Afr. J. Environ. Sci.* 2(1): 85-88.
- Babarinde NAA, Ogunbanjo OO, Babalola JO (2002). Effect of Perming the human scalp hair on its sorption of metal ions. *Afr. J. Technol.* 2(1&2): 200-202
- Babarinde NAA (2002). Adsorption of zinc (II) and cadmium (II) by Coconut husk and goat Hair. *J. Pure Appl. Sci.* 5(1): 81-85.
- Bansode RR, Losso JN, Marshall WE, Rao RM, Portier RJ (2003). Adsorption of metal ions by pecan shell-based granular activated carbons, *Bioresour. Technol.* 89(2): 115–119.
- Cimino G, Passerini A, Toscano G (2000). Removal of toxic cations and Cr (VI) from aqueous solution by hazelnut shell. *Water Res.* 34: 2955–2962.
- Demirbas E (2003). Adsorption of Cobalt (II) from aqueous solution onto activated carbon prepared from hazelnut shells. *Adsorption. Sci. Technol.* 21: 951–963.
- Feng D, Aldrich C, Tan H (2000). Treatment of acid mine water by use of heavy metal precipitation and ion exchange. *Miner. Eng.* 13(6): 623–642.
- Feng D, Aldrich C (2004). Adsorption of heavy metals by biomaterials derived from the marine alga *Ecklonia maxima*. *Hydrometallurgy* 73(1–2): 1–10.
- Fourest E, Roux JC (1992). Heavy metal biosorption by fungal mycelia by-products: Mechanisms and influence of pH. *Appl. Microbiol. Biotech.* 37: 399–403.
- Freundlich H (1907). *Ueber die Adsorption in Loesungen* *Z. Physic. Chem.* 57: 385–470.
- Jain SK, Vasudevan P, Jha NK (1989). Removal of some heavy metals from polluted water by aquatic plants: studies on duckweed and water velvet, *Biol. Wastes.* 28: 115–126.
- Júnior LMB, Macedo GR, Duarte MML, Silva EP, Lobato AKCL (2003). Biosorption of cadmium using the fungus *Aspergillus niger*. *Braz. J. Chem. Eng.* 20(3): 1-17.
- Kobya M, Demirbas E, Senturk E, Ince M (2005). Adsorption of heavy metal ions from aqueous solutions by activated carbon prepared from apricot stone. *Bioresour. Technol.* 96(13): 1518–1512.

- Langmuir I (1918). The adsorption of gases on plane surfaces of glass, mica and platinum. *J. A. Chem. Soc.* 40: 1361–1403.
- Lodeiro L, Cordero B, Barriada JL, Herrero R, Sastre de Vicente ME (2005). Biosorption of cadmium by biomass of brown marine macroalgae. *Bioresour. Technol.* 96(16): 1796-1803.
- Marques PASS, Rosa MF, Pinheiro HM (2000). pH effects on the removal of Cu^{2+} , Cd^{2+} and Pb^{2+} from aqueous solution by waste brewery waste. *Bioprocess Eng.* 23: 135–141.
- Matlock MM, Howerton B, Atwood DA (2002). Chemical precipitation of heavy metals from acid mine drainage. *Water Resources* 36(19): 4757–4764.
- Mohammadi T, Moheb A, Sadrzadeh M Razmi A (2005). Modelling of metal ion removal from waste water by electro dialysis, separate. *Purif. Technol.* 41(1): 73–82.
- Vasuderan P, Padmavathy V, Dhingra SC (2003). Kinetics of biosorption of cadmium on Baker's yeast. *Bioresour. Technol.* 89(3): 281–287.
- Xu H, Liu Y, Tay J (2006). Effect of pH on nickel biosorption by aerobic granular sludge. *Bioresour. Technol.* 97(3): 359–363.